

Protocol cleaning and disinfection

(MS TopFoam LC Fresh – MS Megades Novo)

Cleaning

Step 1: Good preparation and all materials which do not belong there while cleaning, should be removed.

- Remove manure, straw, etc.
- Remove all loose material, such as drinking and feeding troughs where possible
- In case of dirty surfaces, concrete, and / or wooden surfaces, and for the removal of dust, moisten the surface.



Step 2: Foaming of department/room with MS TopFoam LC Fresh.

- Application concentration: 1-2%.
- Please note that all areas where the animal can come into contact with, should be foamed.
- Contact time: minimum 45 minutes up to 120 minutes.
- Do not allow to dry.
- Apply with MS Greenline. (blue side)



Step 3: Rinsing of department/room with high pressure.

Once finished, rinse the barn again briefly (low pressure).



Step 4: Check the department/room visually on quality of cleaning.

Disinfect



Step 5: Let the department/room as possible dry.

- Disinfect with MS Megades Novo / MS Megades Oxy (when under time pressure)
- Please note that all areas where the animal can come into contact with, should be foamed.
- Application concentration: MS Megades Novo 0,75% / MS Megades Oxy 0,5 - 3 %
- Apply with MS Greenline (yellow side)
- Rinsing after contact time (special attention for feeding and drinking trough)
- Let the department/room dry



After about 12 hours, animals can be placed in the section again. If desired, it is possible to take swabs after about 12 hours to control the quality of cleaning and disinfection.

Please note when using MS Greenline:

Approx. 0.4 liters wetting per m² for optimum performance.